

CASE REPORT

BARDET-BIEDL SYNDROME IN AN ETHIOPIAN: A CASE REPORT

Asamere Tsegaw¹, TiliksewTeshome²

ABSTRACT

Bardet-Biedl syndrome (BBS) is a rare familial and multi-system disorder with an autosomal recessive pattern of inheritance and wide range of clinical variability. Its main manifestations are progressive retinal dystrophy, renal dysfunction, post-axial polydactyly, central obesity, mental retardation, and hypogonadism. Renal failure is known to be the main cause of death in patients with BBS. Retinal dystrophy and other eye diseases seen in patients with BBS can cause severe visual impairment and blindness at an early age. We report the clinical and laboratory data of the first published case from Ethiopia of an 18 year old boy with multi-system manifestations of the Bardet-Biedl Syndrome. We discuss the main clinical manifestations of the syndrome including its potentially blinding and fatal features. We emphasize the need for diagnosis of this syndrome at as an early age as possible so that proper and multidisciplinary medical care can be given for such patients to prevent unnecessary morbidity and early mortality.

Key words: Bardet-Biedl syndrome, Retinal dystrophy, polydactyly, renal failure.

INTRODUCTION

The Bardet-Biedl syndrome (BBS) is a familial disorder with an autosomal recessive pattern of inheritance. It is a syndrome with multi-system feature but characterized mainly by progressive retinal photoreceptor cells dystrophy, central obesity, postaxial polydactyly, mental retardation hypogonadism and renal dysfunction. (1,2)

BBS was first described in 1920 by Bardet G. (3) and again in 1922 by Biedl, A. (4). It was often confused with the Laurence-Moon syndrome which was first described in 1866 by Laurence and Moon (5). Laurence and Moon described 4 cases of pigmentary retinal dystrophy, hypogonadism, obesity and spastic paraplegia. These syndromes are similar by being characterized by pigmentary retinopathy, mental retardation, obesity and hypogenitalism. But they differ in that BBS usually includes polydactyly and renal anomalies, whereas the Laurence-Moon syndrome is associated with spastic paraplegia.

Prevalence rates in North America and Europe range from 1:140,000 to 1:160,000 live births. (6) An extensive search of literature revealed no published

case report or large scale case series study from Ethiopia or Africa. To our knowledge, this is the first case report of BBS from Ethiopia.

CASE REPORT

Our patient was an 18 years old boy from Dire Dawa town in the eastern part of Ethiopia. He was seen at a private clinic in Addis Ababa in September 2009 for poor vision. He was then referred to the Retina Clinic at Menelik II Hospital for further evaluation. The chief complaint was progressive reduction of vision in both eyes which was noted since early childhood. His parents admitted that he had delayed developmental milestones like walking and speech when compared to his peers. Although he was currently attending regular school, he repeatedly failed to progress into higher grades and was just a grade 7 student. History of consanguinity was not obtained.

Vital signs were within normal limits. His weight was 93kg with a body mass index (BMI) of 32kg/m². Examination of the abdomen revealed central truncal obesity, (Fig.1) but there was no organomegaly or other abnormalities. Examination of the extremities revealed postaxial extra-fingers or polydactyly (Fig 2, 3) in all upper and lower extremities making the

¹ Department of Ophthalmology, College of Medicine and Health Sciences, The University of Gondar

² Vitreo-Retina unit, Department of Ophthalmology, Faculty of Medicine, Addis Ababa University

* Corresponding Author: Asamere Tsegaw, Email:asameret@yahoo.com

total number of his fingers twenty-four. His fingers, especially in the lower limbs were typically short, broad and stubby (Fig 2). Neurological examination didn't reveal any abnormality.



Figure 1: Central obesity in an 18 year old Ethiopian patient with BardetBiedl syndrome



Figure 2: Sixth toe (polydactyly) in the lower limb of an 18 year old Ethiopian patient with BardetBiedl syndrome



Figure 3: Six fingers (polydactyly) in the upper limbs of an 18 year old Ethiopian patient with BardetBiedl syndrome

Ophthalmic examination showed that his Snellen visual acuity was 1/60 in both eyes. His visual acuity improved to 6/36 in both eyes after refractive correction with -5.0 diopter lens. He also had severe reduction of color vision which was detected on color vision test using Ishihara color plates. Funduscopy examination with the Volk double aspheric +90 diopter lens and fundus camera (Fig 4 and 5) showed that he had signs of bilateral retinal dystrophy whose features included pigmented macular scar, attenuated retinal arterioles, optic nerve head atrophy (“waxy disc pallor”) and patchy retinal pigment clumping. Electrophysiologic tests were not done because of lack of availability.




Figure 4: Fundus picture of the right eye of an 18 year old Ethiopian patient with BardetBiedl syndromeshowing waxy optic disc pallor, myopicperi-papillary atrophy, retinal pigment clumping, and central macular scar



Figure 5: Fundus picture of the left eye of an 18 year old Ethiopian patient with BardetBiedl syndrome showing waxy optic disc pallor, myopicperi-papillary atrophy, retinal pigment clumping, and central macular scar

Laboratory tests and other investigations done included complete blood cell count, ESR, hemoglobin level, VDRL, FBS, chest x-ray and all were within normal limits. Renal function tests revealed abnormal renal function with elevated serum blood urea nitrogen (40mg/dl) and creatinine (3.0mg/dl) as well as proteinuria on urine dipstick analysis. Abdominal ultrasound examination was done for our patient to detect any structural abnormalities in the kidneys, but it revealed no such abnormalities. The patient was referred to Tikur Anbessa Hospital for further evaluation of his renal status. After refractive correction of his myopia and proper documentation of his retinal and other ocular findings, appointment was given him at the Retina Clinic of Menelik II Hospital for further follow up of his eye problems.

DISCUSSION

BBS is genetically heterogeneous and 12 different genes are identified to date whose mutation of each gene alone causes the syndrome. (7, 8) This great genetic heterogeneity is responsible for the observed variable expressivity or wide range of clinical variability.

Most symptoms and signs of the Bardet-Biedl syndrome (BBS) appear after several years of evolution and are not fully apparent during early childhood. Because of the slow development of the clinical features of BBS, the diagnosis remains difficult in young patients. The average age at diagnosis was 9 years in one large population-based survey done in the UK. (2) This is late for such a debilitating condition and our patient was diagnosed at the age of 18 years, much later than the average age in the above report. Considering its adverse prognosis due to its life threatening manifestations, a high index of suspicion is important for early diagnosis.

Central obesity is one of the main manifestations of BBS patients (1, 2). Of the 109 patients in a series reported by Beales P et al., 72% of post pubertal subjects were overweight (BMI >25 kg/m²) of whom 16% had BMI >40 kg/m² which is defined as morbidly obese according to the WHO classification of body mass index. (2) Our patient had a BMI of 32kg/m², within the obese category (BMI >30 kg/m²) of WHO classification. (2)

Renal functional impairment and structural abnormalities, such as renal parenchymal cysts or lobulations, dysplastic kidneys and unilateral agenesis were reported at a varying frequency in the literature (1, 2,

11). Renal functional impairment was recognized as a major feature of BBS phenotype recently. Beales et al in their population based survey used abdominal ultrasound and investigated 57 of the 109 BBS patients involved in their survey and detected one or more of those mentioned renal structural abnormalities in 26(46%) (2). They also found out that six (5%) had chronic renal failure at the time of the survey (2). On the other hand, O'Dea D et al in Canada did a prospective cohort study on 38 confirmed BBS patients over a 6-year period and used both abdominal ultrasound and intravenous pyelogram (IVP) studies on 25 of 38 BBS patients with normal renal function and reported that 96% had some renal structural anomalies (11). Nine (25%) patients, initially with normal renal function, developed renal impairment over the course of the six-year follow up. Eight patients died of renal failure over the course of the follow up period.

Both authors concluded that renal failure is the main cause of death in BBS patients (2, 11). Abdominal ultrasound examination done for our patient didn't reveal any renal structural abnormality, but he had laboratory signs of renal functional impairment which clearly warranted further care and follow up.

Retinal dystrophy is one of the main features of BBS (1, 2). It is occasionally seen in children in the first decade of life, and electroretinogram (ERG) examination can be used to detect it in suspected children with no apparent signs of retinal dystrophy. Signs of retinal dystrophy are seen in all adult patients with BBS (2, 6). Some of the commonest fundus features seen in such patients include a macular scar with or without epiretinal membrane, optic disc atrophy ("waxy" disc pallor), perivascular retinal pigmentary changes in "bonny spicule" pattern or pigment clumping and generalized retinal arterial narrowing (1, 2, 6). Apart from the retinal dystrophy, refractive errors particularly myopia, divergent strabismus and color vision impairment are other ocular features of the syndrome (2, 6). Our patient had all the mentioned ocular features of BBS except strabismus.

Patients with BBS usually have fully formed bony extra-fingers (polydactyly) situated on the lateral border of the hand or foot. It is also one of the major features of BBS, and 70% of the patients can have one or more extra-fingers at birth (1,2). Our patient had extra-fingers in all of his upper and lower extremities.

Hypogonadism, characterized by mal-descended testes, small penis often buried in adipose tissues, is a

universal finding in males and is a major feature of BBS(2). Irregular menstrual cycles may be seen in females (2).

Mental retardation is one of the main features of the syndrome (2), but it is a more disputed feature of BBS as signs of mental retardation are not easily seen. Developmental delay, late in reaching milestones, more specifically delay in walking and speech development and learning difficulties are the other main features manifested in up to half of BBS patients (2).

Clinical features of BBS are divided into primary and secondary features based on the frequency of their appearance in patients (1, 2). The six primary features which were found to appear more frequently are retinal dystrophy, central obesity, polydactyly, mental retardation/learning difficulties, hypogonadism in males and renal dysfunction (1,2). There is also a long list of secondary features (1, 2). To diagnose BBS we need to have at least four of the primary features or three primary features plus two of the secondary features (1, 2). Our patient had five primary features and additional secondary features which were more than enough to fulfill the diagnostic criteria.

After diagnosis, BBS patients need multidisciplinary medical care involving ophthalmologists, nephrologists, and endocrinologists, and base-line investigations which need to be done include blood sugar level, renal function tests, abdominal ultrasound, IVP, urine analysis, refraction and treatment should be started accordingly. The condition should also be explained to parents or patients, and they should also be followed up regularly with frequency based on the severity of their problem.

CONCLUSION

The Bardet-Biedl Syndrome is an autosomal recessive disorder, and it is well known to cause severe visual disability due to retinal dystrophy as well as mortality mainly due to renal failure. Diagnosis of BBS at an early age may help with appropriate medical care to delay the occurrence of some life threatening complications such as renal failure. This can help to improve patients' quality of life and increase their lifespan. Ophthalmologists should be aware of this syndrome, its main ocular and systemic signs as well as the potential systemic complications so that they will be able to diagnose patients with BBS at an earlier age and give appropriate medical care in a multidisciplinary approach involving nephrologists.

REFERENCES

1. Green JS, Parfrey PS, Harnett JD, Farid NR. The cardinal manifestations of Bardet-Biedl syndrome, a form of Laurence-Moon-Biedl syndrome. *N Engl J Med.* 1989 Oct 12; 321 (15):1002-9.
2. Beales PL, Elcioglu N. New criteria for improved diagnosis of Bardet-Biedl syndrome: Results of a population survey. *J Med Genet.* 1999;36:437-446
3. Bardet G. Sur un syndrome d'obésité congénitale avec polydactylie et rétinopathie pigmentaire (contribution à l'étude des formes cliniques de l'obésité hypophysaire). *These de Paris (Le Grand)* 1920; 470: 407.
4. Biedl A. Ein Geschwisterpaar mit adiposogenitaler Dystrophie. *Dtsch Med Wochenschr* 1922; 48: 1630
5. Laurence J Z, Moon R C (1866): Four Cases of ((Retinitis Pigmentosa) occurring in the same Family, and accompanied by general Imperfections of Development. *Ophthalmol Rev.* 1886;2 3241.
6. Alessandro L F. Benedetto. The ocular phenotype of the Bardet-Biedl syndrome Comparison to non-syndromic retinitis pigmentosa. *Retinal Regenerative disease and Experimental Therapy.* 1999; pp 189-199
7. Ansley SJ, Badano JL, Blacque OE. "Basal body dysfunction is a likely cause of pleiotropic Bardet-Biedl syndrome". *Nature* 425 (6958): 628-33
8. Blacque OE, Reardon MJ, Li C. "Loss of C. elegans BBS-7 and BBS-8 protein function results in cilia defects and compromised intraflagellar transport." *Genes Dev.* 18: 1630-42
9. Nicholas Katsanis, James R. Lupski, Philip L. Beales. Exploring the molecular basis of Bardet-Biedl syndrome D- *Human Molecular Genetics*, 2001, Vol. 10, No. 20 2293-2299
10. Kwitek-Black AE, Carmi R, Geoffrey M.: Linkage of Bardet-Biedl syndrome to chromosome 16q and evidence for non-allelic genetic heterogeneity. *Nature Genet* 5:392, 1993
11. O'Dea D, Parfrey PS, Harnett JD, Hefferton D, Cramer BC, Green J. The importance of renal impairment in the natural history of Bardet-Biedl syndrome. *Am J Kidney Dis* 1996;27:776-83